

INSTRUCTIONS FOR USE

for doctor or healthcare professional use only

PHQ-9 QUICK DEPRESSION ASSESSMENT

For initial diagnosis:

1. Patient completes PHQ-9 Quick Depression Assessment on accompanying tear-off pad.
2. If there are at least 4 ✓s in the blue highlighted section (including Questions #1 and #2), consider a depressive disorder. Add score to determine severity.
3. **Consider Major Depressive Disorder**
 - if there are at least 5 ✓s in the blue highlighted section (one of which corresponds to Question #1 or #2).**Consider Other Depressive Disorder**
 - if there are 2 to 4 ✓s in the blue highlighted section (one of which corresponds to Question #1 or #2).

Note: Since the questionnaire relies on patient self-report, all responses should be verified by the clinician, and a definitive diagnosis is made on clinical grounds, taking into account how well the patient understood the questionnaire, as well as other relevant information from the patient. Diagnoses of Major Depressive Disorder or Other Depressive Disorder also require impairment of social, occupational, or other important areas of functioning (Question #10) and ruling out normal bereavement, a history of a Manic Episode (Bipolar Disorder), and a physical disorder, medication, or other drug as the biological cause of the depressive symptoms.

To monitor severity over time for newly diagnosed patients or patients in current treatment for depression:

1. Patients may complete questionnaires at baseline and at regular intervals (eg, every 2 weeks) at home and bring them in at their next appointment for scoring or they may complete the questionnaire during each scheduled appointment.
2. Add up ✓s by column. For every ✓:
“Several days” = 1 “More than half the days” = 2 “Nearly every day” = 3
3. Add together column scores to get a TOTAL score.
4. Refer to accompanying PHQ-9 Scoring Card to interpret the TOTAL score.
5. Results may be included in patients’ files to assist you in setting up a treatment goal, determining degree of response, as well as guiding treatment intervention.

PHQ-9 SCORING CARD FOR SEVERITY DETERMINATION

for healthcare professional use only

Scoring—add up all checked boxes on PHQ-9

For every ✓: Not at all = 0; Several days = 1;
More than half the days = 2; Nearly every day = 3

Interpretation of Total Score

Total Score	Depression Severity
1-4	Minimal depression
5-9	Mild depression
10-14	Moderate depression
15-19	Moderately severe depression
20-27	Severe depression

CUESTIONARIO MÉDICO PARA EL PACIENTE (PHQ-9)

Nombre: _____ Fecha: _____

¿Ha padecido alguno de los trastornos siguientes en las últimas 2 semanas?
(Responda con una "✓")

	No	Varios días	Más de la mitad de los días	Prácticamente todos los días
1. He tenido poco interés o gusto por hacer las cosas	0	1	2	3
2. Me he sentido triste, deprimido o desesperanzado	0	1	2	3
3. He tenido dificultad para conciliar el sueño o dormir, o duermo demasiado	0	1	2	3
4. Me he sentido cansado o desganado	0	1	2	3
5. No tengo apetito o como demasiado	0	1	2	3
6. Me he sentido mal, como que soy un fracaso o que he defraudado a mi familia o a mí mismo	0	1	2	3
7. He tenido dificultad para concentrarme en cosas como leer el periódico o mirar la televisión	0	1	2	3
8. Me muevo o hablo con tanta lentitud que otros lo han notado o por el contrario, estoy tan inquieto y agitado que he estado más activo de lo normal	0	1	2	3
9. He pensado que sería mejor morir o en hacerme daño	0	1	2	3

Sume las columnas:

+ +

TOTAL:

(Al personal médico: Para fines de interpretación del total, consulte las instrucciones de la portada de la almohadilla desprendible)

10. Si padeció alguno de los trastornos, ¿en qué grado le dificultaron su trabajo, la atención del hogar o su convivencia con otros?

Nada _____
Algo _____
Mucho _____
Demasiado _____

PATIENT HEALTH QUESTIONNAIRE (PHQ-9)

Name: _____ Date: _____

Over the last 2 weeks, how often have you been bothered by any of the following problems?
(use "✓" to indicate your answer.)

	Not at all	Several days	More than half the days	Nearly every day
1. Little interest or pleasure in doing things	0	1	2	3
2. Feeling down, depressed, or hopeless	0	1	2	3
3. Trouble falling or staying asleep, or sleeping too much	0	1	2	3
4. Feeling tired or having little energy	0	1	2	3
5. Poor appetite or overeating	0	1	2	3
6. Feeling bad about yourself—or that you are a failure or have let yourself or your family down	0	1	2	3
7. Trouble concentrating on things, such as reading the newspaper or watching television	0	1	2	3
8. Moving or speaking so slowly that other people could have noticed. Or the opposite—being so fidgety or restless that you have been moving around a lot more than usual	0	1	2	3
9. Thoughts that you would be better off dead, or of hurting yourself in some way	0	1	2	3

Add columns: + +

(Healthcare professional: For interpretation of TOTAL, please refer to instructions on tear-off pad cover)

TOTAL:

10. If you checked off *any* problems, how difficult have these problems made it for you to do your work, take care of things at home, or get along with other people?

Not difficult at all _____
Somewhat difficult _____
Very difficult _____
Extremely difficult _____

BACK OF TEAR PAD

Patient Health Questionnaire (PHQ-9) © 1999 Pfizer Inc. All rights reserved. The names PRIME-MD® and PRIME MD TODAY® are registered trademarks of Pfizer Inc. Patient Health Questionnaire (PHQ-9) is adapted from PRIME MD TODAY, developed by Drs. Robert L. Spitzer, Janet B.W. Williams, Kurt Kroenke, and colleagues, with an educational grant from Pfizer Inc.



LM1201161

© 2003 Pfizer Inc.

All rights reserved.

Printed in USA/October 2003

