

References

- ¹Kroenke, K, Spitzer, RL, Williams, JBW (2001). The PHQ-9: Validity of a brief depression severity measure. *Journal of General Medicine*, 16, 606-613.
- ²Spitzer, RL, Kroenke, K, Williams, JBW. (1999). Patient health questionnaire study group. Validity and utility of a self-report version of PRIME-MD: The PHQ primary care study group. *JAMA*, 282, 1737-44.
- ³Spitzer, RL, Williams, JBW, Kroenke, j et al. (2000). Validity and utility of the Patient health Questionnaire in assessment of 3000 obstetric-gynecologic patients: the PRIME-MD Patient Health Questionnaire Obstetrics-Gynecology Study. *American Journal of Obstetrics and Gynecology*, 183:759-69.
- ⁴Patient Health Questionnaire (PHQ-9). <http://www.pfizer.com/pfizer/download/do/phq-9.pdf>. Retrieved May 23, 2007.
- ⁵Patient Health Questionnaire. <http://www.depression-primarycare.org/clinicians/toolkits/materials/forms/phq9/>. Retrieved May 23, 2007.
- ⁶Wulsin, L, Somoza, E & Heck, J (2002). The feasibility of using the Spanish PHQ-9 to screen for depression in primary care in Honduras. *Primary Care Companion Journal Clinical Psychiatry*, 191-195.
- ⁷Bledsoe, SE, & Grote, NK (2006). Treating depression during pregnancy and the postpartum: A preliminary meta-analysis. *Research in Social Work Practice*, 16 (2), 109-120.